



National
Aeronautics and
Space
Administration

NASA Scientific and Technical Document Availability Authorization (DAA)

The DAA approval process applies to all forms of published NASA Scientific and Technical Information (STI), whether disseminated in print or electronically. It is to be initiated by the responsible NASA Project Officer, Technical Monitor, author, or other appropriate NASA official for all presentations, reports, papers, and proceedings that contain NASA STI. Explanations are on the back of this form and are presented in greater detail in NPG 2200.2, "Guidelines for Documentation, Approval, and Dissemination of NASA Scientific and Technical Information."

☐ Original
☐ Modified

I. DOCUMENT/PROJECT IDENTIFICATION

TITLE Protecting the environment in the space age: GIS at the John F. Kennedy Center.	AUTHOR(S) Duncan, B.W., R.A. Reddick, R.B. Smith and D.R. Breininger.
--	--

ORIGINATING NASA ORGANIZATION Kennedy Space Center Biomedical Office	PERFORMING ORGANIZATION (If different) Dynamac Corporation, KSC	
CONTRACT/GRANT/INTERAGENCY/PROJECT NUMBER(S)	DOCUMENT NUMBER(S) Geo Info Systems 4:38-42	DOCUMENT DATE 1994

For presentations, documents, or other STI to be externally published (including through electronic media), enter appropriate information on the intended publication such as name, place, and date of conference, periodical, or journal name, or book title and publisher in the next box. These documents must be routed to the NASA Headquarters or Center Export Control Administrator for approval (see Sections III and VIII).



II. SECURITY CLASSIFICATION

CHECK ONE (One of the five boxes denoting Security Classification must be checked.)

☐ SECRET ☐ SECRET RD ☐ CONFIDENTIAL ☐ CONFIDENTIAL RD ☒ UNCLASSIFIED

III. AVAILABILITY CATEGORY





<input type="checkbox"/> ITAR <input type="checkbox"/> EAR	Export Controlled Document - USML Category /CCL Export Control Classification Number (ECCN) (Documents marked in this block must have the concurrence/approval of the NASA Headquarters or Center Export Control Administrator (see Section VIII).)
<input type="checkbox"/> TRADE SECRET <input type="checkbox"/> SBIR <input type="checkbox"/> COPYRIGHTED	Confidential Commercial Document (check appropriate box at left and indicate below the appropriate limitation and expiration): <input type="checkbox"/> U.S. Government agencies and U.S. Government agency contractors only <input type="checkbox"/> NASA contractors and U.S. Government only <input type="checkbox"/> U.S. Government agencies only <input type="checkbox"/> NASA personnel and NASA contractors only <input type="checkbox"/> NASA personnel only <input type="checkbox"/> Available only with the approval of issuing office: <input type="checkbox"/> Limited until (date)
<input type="checkbox"/> PUBLICLY AVAILABLE	Publicly available documents must be unclassified, may not be export controlled, may not contain trade secret or confidential commercial data, and should have cleared any applicable patents application process.

IV. DOCUMENT DISCLOSING AN INVENTION

THIS DOCUMENT MAY BE RELEASED ON (date)	NASA HQ OR CENTER PATENT OR INTELLECTUAL PROPERTY COUNSEL SIGNATURE	DATE
---	--	-------------

V. BLANKET RELEASE (OPTIONAL)

- ☐ All documents issued under the following contract/grant/project number may be processed as checked in Sections II and III.
- ☐ The blanket release authorization granted on (date)
- ☐ is RESCINDED - Future documents must have individual availability authorizations.
- ☐ is MODIFIED - Limitations for all documents processed in the STI system under the blanket release should be changed to conform to blocks as checked in Sections II and III.

VI. AUTHOR/ORIGINATOR VERIFICATION			
I HAVE DETERMINED THAT THIS PUBLICATION:			
<input type="checkbox"/> DOES contain export controlled, confidential commercial information, and/or discloses an invention for which a patent has been applied, and the appropriate limitation is checked in Sections III and/or IV.			
<input checked="" type="checkbox"/> does NOT contain export controlled, confidential commercial information, nor does it disclose an invention for which a patent has been applied, and may be released as indicated above.			
NAME OF AUTHOR/ORIGINATOR	MAIL CODE	SIGNATURE	DATE
Brean Duncan	Dyn-2		4/6/98
VII. PROJECT OFFICER/TECHNICAL MONITOR/DIVISION CHIEF REVIEW			
<input checked="" type="checkbox"/> APPROVED FOR DISTRIBUTION AS MARKED ON REVERSE <input type="checkbox"/> NOT APPROVED			
NAME OF PROJECT OFFICER OR TECH. MONITOR	MAIL CODE	SIGNATURE	DATE
RAYMOND M WHEELER	JJ-G		4/7/98
VIII. EXPORT CONTROL REVIEW/CONFIRMATION			
<input checked="" type="checkbox"/> Public release is approved <input type="checkbox"/> Export controlled limitation is not applicable			
<input type="checkbox"/> Export controlled limitation is approved <input type="checkbox"/> Export controlled limitation (ITAR/EAR) marked in Section III is assigned to this document:			
USML CATEGORY NUMBER	CCL ECCN NUMBER	HQ OR CENTER EXPORT CONTROL ADMINISTRATOR (as applicable)	DATE
			4/16/98
IX. PROGRAM OFFICE OR DELEGATED AUTHORITY REVIEW			
<input checked="" type="checkbox"/> APPROVED FOR DISTRIBUTION AS MARKED ON REVERSE <input type="checkbox"/> NOT APPROVED CODE M, DAA LETTER, 6/14/94			
NAME OF PROGRAM OFFICE REPRESENTATIVE	MAIL CODE	SIGNATURE	DATE
Audrey Lee Silipo	FF-S2-A		4/14/98
X. DISPOSITION			
THIS FORM, WHEN COMPLETED, IS TO BE SENT TO YOUR CENTER PUBLICATIONS OFFICE			

INSTRUCTIONS FOR COMPLETING THE NASA SCIENTIFIC AND TECHNICAL DOCUMENT AVAILABILITY AUTHORIZATION (DAA) FORM

Purpose. This DAA form is used to prescribe the availability and distribution of all NASA-generated and NASA-funded documents containing scientific and technical information (including those distributed via electronic media such as the World Wide Web and CD-ROM).

Requirements. The author/originator must provide either a suitable summary description (title, abstract, etc.) or a completed copy of the document with this form. This form is initiated by the document author/originator and that individual is responsible for recommending/determining the availability/distribution of the document. The author/originator completes Sections I through III, and VI. The author/originator is also responsible for obtaining information and signature in Section IV to the extent the document discloses an invention for which patent protection has been applied. Subsequent to completion of these sections, the author/originator forwards the document to the appropriate Project Manager/Technical Monitor/Division Chief for further review and approval in Section VII, including a re-review of the planned availability and distribution. Once this approval is obtained, the DAA is forwarded to the NASA Headquarters or Center Export Administrator for completion of Section VIII. It is then forwarded for completion of Section IX to the cognizant NASA Headquarters Program Office or Delegated Authority, who provides final review and approval for release of the document as marked.

When to Use This Form. Documents containing STI and intended for presentation or publication (including via electronic media) must be approved in accordance with the NASA STI Procedures and Guidelines (NPG 2200.2). Documents that are to be published in the NASA STI Report Series must be coordinated with the appropriate NASA Headquarters or Center Scientific and Technical Information Office in accordance with NPG 2200.2. Note that information on the Report Documentation Page (if attached) is not to be entered on the DAA except for title, document date, and contract number.

How to Use this Form. Specific guidelines for each section of this form are detailed below.

I. Document/Project Identification. Provide the information requested. If the document is classified, provide instead the security classification of the title and abstract. (Classified information must not be entered on this form). Include RTOP numbers on the Contract/Grant/Interagency/Project Number(s) line. Provide information on presentations or externally published documents as applicable.

II. Security Classification. Enter the applicable security classification for the document. Documents, if classified, will be available only to appropriately cleared personnel having a "need to know."

III. Availability Category for Unclassified Documents. Check the appropriate category or categories.

Export Controlled Document. If the document is subject to export restrictions (see NPG 2200.2, paragraph 4.5.3), the appropriate restriction must be checked, either International Traffic in Arms Regulations (ITAR) or Export Administration Regulations (EAR), and the appropriate United States Munitions List (USML) category or Commerce Control List (CCL), Export Control Classification Number (ECCN) must be cited.

Confidential Commercial Documents (Documents containing Trade Secrets, SBIR documents, and/or Copyrighted Information). Check the applicable box (see NPG 2200.2 paragraph 4.5.7). When any of these boxes are checked, also indicate the appropriate limitation and expiration in the list to the right of these restrictions. These limitations refer to the user groups authorized to obtain the document. The limitations apply both to the initial distribution of the documents and the handling of requests for the documents. The limitations will appear on and apply to reproduced copies of the document. Documents limited to NASA personnel should not be made available to onsite contractors. If the Available Only With the Approval of Issuing Office limitation is checked, the NASA Center for Aerospace Information will provide only bibliographic processing and no initial distribution; CASI will refer all document requests to the issuing office.

Publicly Available Document - Unrestricted Distribution. Check this box if the information in the document may be made available to the general public without restrictions (unrestricted domestic and international distribution). If the document is copyrighted (see paragraph 4.5.7.3 in NPG 2200.2), also check the "Copyrighted" box in this section.

IV. Document Disclosing an Invention. This must be completed when the document contains information that discloses an invention (see NPG 2200.2, paragraph 4.5.9). When this box is checked, an additional appropriate availability category must be checked. Use of this category must be approved by NASA Headquarters or Center Patent Counsel or the Intellectual Property Counsel.

V. Blanket Release (Optional). Complete this optional section whenever subsequent documents produced under the contract, grant, or project are to be given the same distribution and/or availability as described in Sections II and III. More than one contract number or RTOP Number can be entered. This section may also be used to rescind or modify an earlier Blanket Release. All blanket releases must be approved by the Program Office or its designee and concurred with by the Office of Management Systems and Facilities.

VI. Author/Originator Verification. Required for all DAA forms.

VII. Project Officer/Technical Monitor/Division Chief Review. The Project Officer/Technical Monitor/Author or Originator Division Chief or above must sign and date the form. The office code and typed name should be entered.

VIII. Export Control Review/Confirmation. This section is to be completed by the authorized NASA Headquarters or Center Export Control Administrator for all documents.

IX. Program Office or Delegated Authority Review. This section is to be completed by the duly authorized official representing the NASA Headquarters Program Office. Any delegation from NASA Headquarters to a NASA Center in accordance with NPG 2200.2 should be entered here.

X. Disposition. For NASA Center use.

Protecting the Environment in the Space Age: GIS at the John F. Kennedy Space Center

Brean W. Duncan, Resa A. Reddick, Rebecca B. Smith, and David R. Breininger

The Bionetics Corporation




Figure 1. Kennedy Space Center is located on the central east coast of Florida.

Figure 2. Listed as a federally threatened species by the U.S. Fish and Wildlife Service, a banded Florida scrub jay is shown in preferred habitat with sand openings.



The Biomedical Operations and Research Office (MD) of the National Aeronautics and Space Administration (NASA) is using geographic information systems (GIS) technology in its effort to monitor protected wildlife at Kennedy Space Center (KSC). Data bases containing environmental variables and demographic data have been compiled for a number of threatened and endangered species on KSC, along with data bases showing the effects of Space Shuttle launches on KSC. NASA also uses GIS technology to monitor such episodic environmental impacts as Space Shuttle launch exhaust deposition.

Increased urbanization has reduced the amount and quality of natural habitat in Florida. As urbanization increases, so does the importance of protected lands in the state. Kennedy Space Center (KSC) is the largest protected area on the east coast of Florida and is home to 14 federally listed and 15 state-listed wildlife species identified as threatened or endangered. The National Aeronautics and Space Administration (NASA) has particularly challenging management goals that include using the land for space operations and preserving the valuable habitat contained within KSC. Geographic information systems (GIS) technology is one of the many tools being used to meet those management goals by mapping and monitoring environmental variables and long-term demographic data sets on selected animal species.

KSC, the principal launch site for NASA space systems, is centrally located on the east coast of Florida (Figure 1), where the initial trajectories of space transport systems are over the Atlantic Ocean. NASA began acquiring land in early 1962

on north Merritt Island, Florida, as a for launch operations in support of Manned Lunar Landing Program. now encompasses 57,000 hectare 140,000 acres — which is consid extensive enough to provide surrou communities protection from la operations.

KSC is long and narrow, borders the Indian River Lagoon on the and Cape Canaveral Air Force St (CCAFS) and the Atlantic Ocean o east. Merritt Island and Cape Canaver a barrier island complex of Pleistocen Recent ages. The topography consist series of ridges and swales derived relict dunes deposited as the barrier is were formed. Elevation ranges fror level to as much as nine meters o most recent dunes.

Because KSC was found to inc prime habitat for unique and endan flora and fauna, NASA entered in agreement with the U.S. Fish and Wi Service (USFWS) to establish a wi preserve, known as the Merritt Is National Wildlife Refuge (MINWR), in the boundaries of KSC. A similar a ment was reached with the Departme the Interior to form the Canaveral Nat Seashore (CNS), managed by the Nat Park Service (NPS), within KSC's r ern boundaries. The majority of KSC is undeveloped, with only five pe being used for NASA operations industrial use. The remaining 95 per managed by USFWS and NPS.

KSC is located in a transition between the warm temperate Carol Zoogeographic Province to the north the subtropical Caribbean Zoogeogr Province to the south. The result o location is high biodiversity with sp from each region being present. The more than 1,500 taxa — or biolo groupings — on KSC, including species of fish, 315 species of birc species of amphibians and reptile

Brean W. Duncan is a physical geogr and ecologist, Resa A. Reddick is a cputer/GIS specialist and both Rebecca Smith and David Breininger are wildl biologists. All the authors are with th netics Corporation, NASA Biomedica Operations and Research Office, Ken Space Center, Florida, USA.

cies of mammals, and almost 1,000 species of vascular plants (NASA, 1990; NASA, 1992).

The Biomedical Operations and Research Office (MD) of NASA has been reporting environmental monitoring and research on KSC since the early 1970s. Program elements include monitoring of baseline conditions to document natural variability in the ecosystem, assessments of operations, assessments of new facilities construction, and ecological research using on wildlife habitat associations. The ecological program's objective is to develop information necessary to define and predict environmental problems and factors that may result from activities at KSC. To support these requirements, the ecological program uses a proactive, long-term, multidisciplinary approach (Hinkle et al., 1988).

Remote sensing and GIS technologies are at the center of the environmental monitoring and research activities on KSC. ARC/INFO (Environmental Systems Research Institute [ESRI], Redlands, California, USA) software is used for analyzing both episodic and continuous environmental impacts, as well as threatened and endangered species' demographic-habitat relationships. This process enhances our predictive capability to minimize or mitigate potential environmental impacts on KSC.

APPLICATIONS

Habitat bases containing environmental variables and demographic data have been compiled for a number of threatened and endangered species on KSC. We have focused our energy and resources on species with habitats having the highest probability of being affected by NASA operations and those that are indicators for other species of special concern. Data series showing the effects of Space Shuttle launches on KSC have also been produced. Following are examples of these data series.

Florida scrub jay. Field biology, remote sensing, and GIS methods are combined to map habitat features important for one of the most habitat-specific birds in North America. The Florida scrub jay (*Aphelococcyx coerulescens*), shown in Figure 2, is a threatened species restricted to oak-dominated scrub, which occurs primarily on well-drained soils in the central




Figure 3. Scrub jay source and sink areas created from overlaying 1989–1991 yearling production and breeder mortality coverages for the Tel-4 long-term study area

Florida peninsula. These birds live in territories defended year-round by a permanently monogamous breeding pair and young scrub jays called helpers (Woollenden and Fitzpatrick, 1984).

Urbanization of well-drained ridges throughout Florida has caused a major decline in suitable scrub jay habitat. The result of this habitat destruction and degradation is that KSC is now home to one of the largest remaining Florida scrub jay populations (Breininger, 1989). Proper management of this habitat is critical for the scrub jay's survival.

A data base containing both environmental variables and long-term demographic data has been produced for two study areas on KSC. Habitat maps have been produced using very large-scale, false-color aerial imagery and a detailed Anderson Land-use/Landcover classification system. Each scrub jay in the study areas has been banded using a unique combination of two-color bands and an aluminum USFWS band. Territory boundaries were mapped during April and May using repeated visits to determine the area defended by each family. Nearly all the nest attempts within the study boundaries



Figure 4. The gopher tortoise is listed as a species of special concern by the Florida Game and Fresh Water Fish Commission.




Figure 5. Each gopher tortoise home-range polygon was created by overlaying all burrows used by individual tortoises and connecting the outermost burrows. Solid and dashed arcs represent male and female tortoise home ranges, respectively, and are shown here overlaid on habitat polygons.

during each nesting season were located and mapped. These maps were digitized into separate coverages representing each season, and demographic attributes were stored along with their corresponding spatial locations and areas.

Territories, nest locations, and demographic data were overlaid on the habitat map and other environmental layers to gain further insight into the scrub jay's habitat requirements. Some of the analyses were as simple as calculating arithmetic means of territory sizes and acreages of habitat types within territories for each year. Some analyses involved demographic

data entirely, such as differencing breeder mortality and yearling production to produce coverages of source (areas where yearling production exceeds breeder mortality) and sink (areas where breeder mortality exceeds yearling production) locations, as shown in Figure 3. Other, more analytical techniques involved determining correlations and associations between habitat and demographic variables. These studies allow the development and testing of habitat models for environmental assessments and mitigation of environmental impacts on KSC. GIS techniques enhance the mapping and delineation of the most important areas for maintaining a viable Florida scrub jay population on KSC, thereby enhancing land-use planning activities (Breininger, et al, 1991).

The gopher tortoise (*Gopherus polyphemus*) is a 3.5- to 4.5-kilogram animal, shown in Figure 4, that can be found from extreme southern South Carolina and eastern Louisiana through much of mainland Florida. The tortoises occupy a variety of habitats but require areas with an adequate cover of herbaceous plants for food and suitable soils in which to dig burrows (Cox et al., 1987). Burrows provide protection from fires, predators, and harsh climatic conditions. In addition, at least 39 invertebrate and 42 vertebrate species have been documented using gopher tortoise burrows. A combination of habitat destruction, predation by humans and other animals, and unique biological characteristics have placed the gopher tortoise's continued existence in jeopardy. It is listed as a "species of special concern" by the Florida Game and Fresh Water Fish Commission (Wood, 1986).

KSC is the largest area of protected habitat for the gopher tortoise on the Atlantic coast of Florida (Breininger et al., 1988). A radio telemetry study was done at two sites on KSC between 1987 and 1989 to answer questions about habitat and burrow use. Twenty-three tortoises were trapped, outfitted with transmitters, and tracked twice per week for an average of 10 months. Burrow locations for the tagged tortoises were digitized from color infrared aerial photographs into the GIS. Because a single tortoise uses several burrows within an area, home-range polygons were developed for each animal based on the burrow locations. Individual home-range sizes were determined, as well as




Figure 6. The West Indian manatee is listed as an endangered species by the U.S. Fish and Wildlife Service.

average home range sizes for males and females.

The home range polygons were overlaid onto vegetation coverages to examine the types and amounts of habitats within them, as shown in Figure 5. These data, when considered with other information — for example, the number of burrows used per tortoise, as well as seasonal activity — provide a visual representation of tortoise habitat use that cannot be achieved with numbers alone. Results can be used to estimate population sizes and determine critical habitat requirements, both of which are important considerations for a species so greatly influenced by land management practices.

The West Indian manatee (*Trichechus manatus*), shown in Figure 6, sometimes called the Florida manatee, was federally listed as an endangered species in 1972. Manatees are basically estuarine in Florida, although they are occasionally seen along the beaches. Manatees are migratory and can range in excess of 820 kilometers (Reid, 1991). They are found throughout Florida and southeastern Georgia and occasionally along coastal states farther north and west.

Manatees have always been found in the surrounding waters of KSC, with the highest peak during spring and a smaller surge during fall. In 13 years of monitoring, a five-fold increase of manatees using KSC waters during spring has been documented. The spring peak in recent years has includ-



Figure 7. Summer manatee sightings for 1977 through 1991 overlaid on a Landsat scene. Clustered sightings indicate preferred habitat locations during summer season.

ed more than 300 manatees within the KSC boundaries of the Banana River (Provancha and Provancha, 1988). The increased use of KSC waters by manatee appears to manifest a shift away from previously used areas and reflect the degradation of habitat outside protected areas. Manatee mortalities have increased at an alarming rate in Florida, with human development and boating activities being the main causes (Beck et al., 1982).




Figure 8. A Space Shuttle blasts off from Kennedy Space Center, causing the formation and dispersion of an exhaust cloud.




Figure 9. Cumulative Space Shuttle near-field exhaust deposition pattern for Pad 39A. Frequent deposition events occur close to the pad and decrease with distance.

All open water on KSC deeper than one meter is potential habitat for the manatee, extensive seagrass beds bordering deep bays are preferred. Fresh water sources to attract manatees; some researchers believe fresh water to be a requirement. Since 1977, aerial surveys have been conducted over KSC waters to monitor the

number and distribution of manatees. Surveys are conducted with light planes or helicopters using a set flight path. During the early surveys, observers would call out sightings and any behavior associated with them and a person would mark the data on a map. These maps were stored in a file drawer until 1989, when they were digitized into ERDAS (Atlanta, Georgia, USA) image processing software and converted into ARC/INFO using GENERATE to produce a point coverage. Problems with this procedure such as positional accuracy, map accuracy and points being omitted during digitization have forced more automated techniques to be used. Since 1990, an Apollo II

IV (Ashton-Tate, Torrance, California) into ASCII format with latitude and longitude conversions to State Plane coordinates performed in ERDAS. The data are then loaded into ARC/INFO via the GENERATE and ADD FROM commands. This technique is much easier and more time efficient than its predecessor.

Our GIS has provided a means for storage and analysis of a large, long-term, and comprehensive manatee data base (Figure 7). The GIS data allow NASA and the USFWS to track manatee time and space interactions, which are important for performing facilities assessments on KSC. By using this valuable information, NASA can manage KSC activities to minimize impact on the manatee's fragile habitat.

LAUNCH DEPOSITION MONITORING

NASA also uses GIS technology to monitor such episodic environmental impacts as Space Shuttle launch exhaust deposition. Space Shuttle launches produce localized acidic deposition through the formation and dispersion of an exhaust cloud, shown in Figure 8. The cloud contains carbon dioxide, aluminum oxide, and hydrogen chloride, along with additional minor components. These cloud constituents, when combined with water vapor and 300,000 gallons of sound suppression and cooling water dumped on the pad at the time of launch, produce acidic deposition (Anderson and Keller, 1983). Typically this cloud is directed northward by a flame trench and then rises to a stabilization height and is carried by prevailing winds.

Two divisions of deposition exist: near-field and far-field. Near-field deposition occurs on the immediate north side of the launch pad and involves the highest concentration of acid. Far-field deposition occurs after the cloud rises and moves with prevailing winds. Areas receiving far-field deposition, which is primarily dry residue, vary with meteorological conditions and occur farther away from the pad. Both near- and far-field deposition are mapped by field survey after each launch event.

The near-field and far-field maps from each launch are digitized into the GIS (for example, Duncan and Schmalzer, 1992). Individual maps of the deposition are combined to produce cumulative deposition patterns for both Shuttle launch facilities (Figures 9 and 10). Three different maps

are continually updated, near-field maps for launch pads 39A and 39B and a combined far-field map for both pads.

Cumulative maps of deposition reveal the frequency of deposition at any given location on KSC from the 11 years of Shuttle launches, as well as which launches are responsible for that deposition. The data base can be updated after each launch, producing a continuous source of spatial information to determine and monitor the frequency of cumulative deposition, which has implications for both operational uses and habitat management for the flora and fauna of KSC.

CONCLUSION

NASA is committed to protecting the environment of KSC while continuing to launch space transport systems. Proper management and monitoring of this protected land is increasingly important and must be based on sound scientific data. GIS methods have enhanced NASA's ability to monitor and manage natural resources on KSC by providing an efficient means of data storage and spatial-numerical analysis. The ongoing long-term studies cur-

rently underway on KSC produce large quantities of spatial data and even larger quantities of relational data.

GIS technology is increasing the ability of scientists on KSC to analyze their data in a spatially explicit manner and gain greater insight into spatial, temporal, habitat, and animal interactions necessary for proper management. By doing so, we are moving in a direction to produce spatial predictive models that will aid in impact assessment and land mitigation practices necessary for cohabitation of wildlife with ongoing space operations.

REFERENCES

- Anderson, B.J., and V.W. Keller. 1983. Space Shuttle Exhaust Cloud Properties. NASA TP 2258.
- Beck, C.A., R.K. Bonde, and G.B. Rathbun. 1982. Analysis of Propeller Wounds on Manatees in Florida. *Journal of Wildlife Management* 46(2):531-535.
- Breining, D.R. 1989. A New Population Estimate for the Florida Scrub Jay on Merritt Island National Wildlife Refuge. *Florida Field Naturalist* 17:25-31.
- Breining, D.R., P.A. Schmalzer, D.A. Rydene, and C.R. Hinkle. 1988. *Burrow and Habitat Relationships of the Gopher Tortoise in Coastal Scrub and Slash Pine Flatwoods on Merritt Island, Florida*. Florida Game and Fresh Water Fish Commission Nongame Wildlife Program Final Report. Tallahassee, Florida: Florida Game and Fresh Water Fish Commission.
- Breining, D.R., M.J. Provancha, and R.B. Smith. 1991. Mapping Scrub Jay Habitat for Purposes of Land-Use Management. *Photogrammetric Engineering and Remote Sensing* 57(11):1467-1474.
- Cox, J., D. Inkley, and R. Kautz. 1987. Ecology and Habitat Protection Needs of the Gopher Tortoise (*Gopherus polyphemus*) Populations Found on Lands Slated for Large-Scale Development in Florida. Florida Game and Fresh Water Fish Commission Nongame Wildlife Program Technical Report No. 4. Tallahassee, Florida: Florida Game and Fresh Water Fish Commission.
- Duncan, B.W., and P.A. Schmalzer. 1992. "Using ARC/INFO to Monitor Space Shuttle Launch Effects and Validate a Spatial Predictive

Model." In *Proceedings of the Twelfth Annual ESRI User Conference*. Redlands, California: Environmental Systems Research Institute.

EG&G Florida, and Edward E. Clark Engineers-Scientists. 1992. Environmental Resources Document, John F. Kennedy Space Center, NASA, KSC-DF-3080.

Hinkle, C.R., C.R. Hall, M.J. Provancha, P.A. Schmalzer, A.M. Koller, Jr., and W.M. Knott, III. 1988. *The Use of Remotely Sensed Environmental Data and Geographic Referenced Data Management for Environmental Impact Assessments*. Chemical Propulsion Information Agency Publication No. 485.

Provancha, J.A., and M.J. Provancha. 1988. Long-Term Trends in Abundance and Distribution of Manatees (*Trichechus manatus*) in the Northern Banana River, Brevard County, Florida. *Marine Mammal Science* 4(4):323-338.

Reid, J.P., G.B. Rathbun. 1991. Distribution Patterns of Individually Identifiable West Indian Manatees (*Trichechus manatus*) in Florida. *Marine Mammal Science* 7(2):180-190.

Schmalzer, P.A., and C.R. Hinkle. 1990. *Flora and Threatened and Endangered Plants of John F. Kennedy Space Center, Florida*. NASA TM 102791.

Wood, D. 1986. *Official Lists of Endangered and Potentially Endangered Fauna and Flora in Florida*. Tallahassee, Florida: Florida Game and Fresh Water Fish Commission.

Woolfenden, G.E., and J.W. Fitzpatrick. 1984. *The Florida Scrub Jay: Demography of a Cooperative-Breeding Bird*. Monograph in Population Biology No. 20. Princeton, New Jersey: Princeton.

These studies have been conducted under NASA contracts NAS10-10285 and NAS-11624. The authors would like to thank Dr. William Knott, III, and Burton Summerfield, NASA Biomedical Operations and Research Office and Bionetics Program Managers Carlton Hall and Ross Hinkle for their assistance and support. We would also like to thank Mark Provancha, Jane Provancha, Vickie Larson, and Dr. Paul Schmalzer for contributing a considerable amount of work and expertise to these studies.




Figure 10. Cumulative Space Shuttle far-field exhaust deposition for both pads 39A and 39B. Far-field deposition occurs over a much larger area than near-field deposition but occurs less frequently at any one location.